

Planck mission*: challenges and expectations for cosmology and particle physics

Carlo Baccigalupi, SISSA

*After about 16 years of preparation, launching on May 14th, 2009

Outline


- CMB physics
- Status of CMB observations
- Challenges for future CMB
- The Planck mission
- Conclusions

CMB physics



CMB: where, when and how



- Opacity: $\lambda = (n_e \sigma_T)^{-1} \ll H^{-1}$
- Decoupling: $\lambda \approx H^{-1}$
- Free streaming: $\lambda \gg H^{-1}$
- Cosmological expansion, constants and baryon abundance conspire to activate decoupling about 300000 years after the Big Bang, at about 3000 K photon temperature
- Expansion and the metric perturbations affect all cosmological species
- The CMB is a snapshot of cosmological perturbations in the photon component only




CMB: where, when and how

B99

- Opacity: $\lambda = (n_e \sigma_T)^{-1} \ll H^{-1}$
- Decoupling: $\lambda \approx H^{-1}$
- Free streaming: $\lambda \gg H^{-1}$
- Cosmological expansion, constants and baryon abundance conspire to activate decoupling about 300000 years after the Big Bang, at about 3000 K photon temperature
- Expansion and the metric perturbations affect all cosmological species
- The CMB is a snapshot of cosmological perturbations in the photon component only






COsmic Background Explorer (COBE)



Nobel prize for physics, John Mather e George Smoot, 2006

Wilkinson Microwave Anisotropy Probe (WMAP)



CMB physics: Boltzmann equation

$$\frac{d \text{ photons}}{dt} = \text{metric} + \text{Compton scattering}$$

$$\frac{d \text{ baryons+leptons}}{dt} = \text{metric} + \text{Compton scattering}$$

CMB physics: Boltzmann equation

d neutrinos






$$\frac{d \text{ neutrinos}}{dt} = \text{metric} + \text{weak interaction}$$

d dark matter

$$\frac{d \text{ dark matter}}{dt} = \text{metric} + \text{weak interaction (?)}$$



metric = photons + neutrinos + baryons + leptons + dark matter

CMB physics: metric




CMB Physics: Compton scattering

- Compton scattering is anisotropic
- An anisotropic incident intensity determines a linear polarization in the outgoing radiation
- At decoupling that happens due to the finite width of last scattering and the cosmological local quadrupole




CMB anisotropy: total intensity





CMB anisotropy: polarization


Curl (B):




Gradient (E):



CMB anisotropy: reionization




CMB anisotropy: lensing




CMB anisotropy: lensing

E
B



CMB anisotropy: lensing

E
B



Status of CMB observations



CMB anisotropies

$T(\hat{n}), Q(\hat{n}), U(\hat{n}), V(\hat{n})$


$a_{lm}^T, a_{lm}^E, a_{lm}^B$

spherical
harmonics

information
compression

$$C_l = \sum_m (a_{lm}^{T,E,B})(a_{lm}^{T,E,B})^* / 2(l+1)$$

CMB angular power spectrum



Angle $\approx 200/\ell$ degrees


CMB angular power spectrum

Acoustic oscillations

Primordial power

Reionization


Lensing



Angle $\approx 200/l$ degrees


Gravity waves

WMAP first year







Angle $\approx 200/l$ degrees

WMAP fifth year




Angle $\approx 200/\ell$ degrees



CMB angular power spectrum






Cosmological concordance model



Cosmological concordance model





Cosmological concordance model



CMB anisotropy statistics: unknown, probably still hidden by systematics

- Evidence for North south asymmetry (Hansen et al. 2005)
- Evidence for Bianchi models (Jaffe et al. 2006)
- Poor constraints on inflation, the error is about 100 times the predicted deviations from Gaussianity (Komatsu et al. 2003)
- Lensing detection out of reach or marginal, see smith et al. for a 3.4σ detection correlating WMAP and NVSS galaxies





Other cosmological backgrounds?

- Neutrinos: abundance comparable to photons ☺, decoupling at MeV ☺, cold as photons ☹, weak interaction ☹
- Gravity waves: decoupling at Planck energy ☺, abundance unknown ☹, gravitational interaction ☹
- Morale: insist with the CMB, still for many years...that's the best we have for long...

Forthcoming CMB polarization probes



- Planck
- B modes from GWs: EBEx (US, collaborators in France, Italy, UK), balloon, same launch time scale as Planck for the north american flight, SPIDER (US, ...), QUIET (US, UK), ground based, Clover (UK, ...), Brain,
- Lensing: ALMA,
- ...
- Complete list available at lambda.gsfc.nasa.gov
- Time scale: approximately one year for test launches



Cosmic vision beyond Einstein

- NASA and ESA put out separate calls of opportunity for a polarization oriented future (2020, 2030) CMB satellite
- Technologies, design, options for joint or separate missions are in proposals which have been submitted in these weeks
- Promises: gravity waves, lensing and high redshift dark energy, inflationary non-Gaussianity

Cosmic vision program logo



Beyond Einstein logo



Challenges for future CMB



Challenges for future CMB


- The sensitivity can be increases with the detector number 😊
- The systematics from the instrument must be controlled at the level of the signal 😞
- The emission from foregrounds may cover the B signal over the all sky, at all frequency 😞

Jarosik et al. 2006



Challenges for future CMB: systematics from beam shape


- Asymmetric beams cause unwanted polarization from total intensity, leakage of E modes into B, ...
- No way to circularize the beams, rather the beam shape has to be reconstructed in flight to subtract the bias from the signal



Challenges for future CMB: foreground emission

Bennett et al. 2006



- In total intensity, at frequencies between 60 and 90 GHz, after cutting out the brightest part of the Galactic emission, the sky is dominated by CMB



Challenges for future CMB: foreground emission

Bennett et al. 2006

- In total intensity, at frequencies between 60 and 90 GHz, after cutting out the brightest part of the Galactic emission, the sky is dominated by CMB
- In polarization, at frequencies between 60 and 90 GHz, after cutting out the brightest part of the Galactic emission, the sky is dominated by CMB





Page et al. 2006


Challenges for future CMB: foreground emission

Bennett et al. 2006


- In total intensity, at frequencies between 60 and 90 GHz, after cutting out the brightest part of the Galactic emission, the sky is dominated by CMB polarization, at frequencies between 60 and 90 GHz, after cutting out the brightest part of the Galactic emission, the sky is dominated by CMB



Challenges for future CMB: foreground emission



Page et al. 2006





Planck reference sky


Are there foreground clean regions at all in polarization?

Page et al. 2006

- WMAP has no detection in large sky areas in polarization
- Very naive estimates may be attempted in those areas, indicating that the foreground level might be comparable to the cosmological B mode at all frequencies, in all sky regions




Living with foregrounds: component separation



$$\mathbf{x}_1 = a_{11} s_1 + a_{12} s_2 + n_1$$

$$\mathbf{x}_2 = a_{21} s_1 + a_{22} s_2 + n_2$$


Living with foregrounds: component separation



$$\mathbf{x}_1 = a_{11} \mathbf{s}_1 + a_{12} \mathbf{s}_2 + \mathbf{n}_1$$

$$\mathbf{x}_2 = a_{21} \mathbf{s}_1 + a_{22} \mathbf{s}_2 + \mathbf{n}_2$$

Living with foregrounds: component separation



$$x = As + n$$

Invert for s !




Living with foregrounds: component separation

$$\mathbf{x} = \mathbf{A}\mathbf{s} + \mathbf{n}$$

- Non-blind approach: use prior knowledge on \mathbf{A} and \mathbf{s} in order to stabilize the inversion, likely to be suitable for total intensity
- Blind approach: do not assume any prior either on \mathbf{A} or \mathbf{s} , likely to be used in polarization
- Parametrization: introduce extra “cosmological parameters” parametrizing the foreground unknowns, and fit the data with those in, marginalizing afterwards, promising results in total intensity, to be tested in polarization
- Relevant literature from Brandt et al. 1994, to Maino et al. 2006, successful applications to COBE, BEAST, WMAP

Component separation in polarization

- Component separation studies how to separate CMB and foregrounds in astrophysical multi-frequency observations
- The independent component analysis exploits the statistical differences between the almost Gaussian CMB and the strongly non-Gaussian foregrounds
- Results are encouraging, although obtained so far without instrumental systematics



Stivoli et al. 2006

The science goals of the Planck satellite

Source: Planck scientific program bluebook,
available at www.rssd.esa.int/Planck

A decorative graphic consisting of several sets of concentric circles, resembling ripples in water, located in the bottom right corner of the slide. The circles are light blue and vary in size and opacity, creating a subtle background element.

Planck

- Launch: May 14th, 2009, Kourou, French Guiana
- Hardware: third generation CMB probe, ESA medium size mission, NASA (JPL, Pasadena) contribution
- Data analysis Software from about 400 collaborators, EU and US
- Two data processing centers (DPCs): Paris + Cambridge (IaP + IoA), Trieste (OAT + SISSA)






Planck contributors



Planck simulations



Cambridge


Paris

Trieste

Planck data processing sites


The DPC

- OAT and SISSA form the Trieste Data Processing Center
- People: three staff, two temporary researchers, post-docs and PhD students at SISSA, two staff, several contract scientists and technologists at OAT
- Processing raw data for all Low Frequency Instrument radiometers (in parallel to the analysis of the bolometers of the High Frequency Instrument in Paris)
- Two parallel machines (ENT at OAT, 256 CPUs, HG1 at SISSA, 160 CPUs, hundreds of Gb RAM, tens of Tb disk space)




The DPC

- Analysis in parallel to the HFI DPC starting from maps, up to the delivery of all the final proprietary scientific deliverables
- May 14, 2009, Planck launch
- Summer 2009, first survey begins
- Summer 2009 + 14 months: first survey ends
- Proprietary period, analysis for two years, final results and publications by the end of 2011, 2012
- Proposal for mission extension being submitted







Planck data

- All sky maps in total intensity and polarization, at 9 frequencies between 30 and 857 GHz
- Angular resolution from 33' to 7' between 30 and 143 GHz, 5' at higher frequencies
- S/N ≈ 10 for CMB in total intensity, per resolution element
- Catalogues with tens of thousands of extra-Galactic sources







PLANCK GALAXY SURVEYS					
	FREQUENCY [GHz]				
	143	217	353	550	850
Confusion limit [mJy, 3σ]	6.3	14.1	44.7	112	251
Planck All Sky Survey sensitivity [mJy, 3σ]	26	37	75	180	300
Planck Deep Survey sensitivity [mJy, 3σ]	10	18.4	49	170	280
Number of galaxies [all sky]	570	860	1700	4400	35000


Planck scientific deliverables: CMB total intensity and the era of imaging




Planck scientific deliverables: CMB polarization



Planck and polarization CMB B modes



Planck scientific deliverables: cosmological parameters



Planck scientific deliverables: inputs for fundamental interactions

- Absolute neutrino mass measure down to fractions of eV
- Upper limit to cosmological gravitational waves corresponding to 10% of the density fluctuation amplitude
- Constraints on Jordan-Brans-Dicke gravity modifications on cosmological scales $\omega_{\text{JBD}} > 3000$
- Constraints on primordial non-Gaussianity ten times better than present constraints, unprecedented foreground control
- Overall statistical assessment of CMB anisotropy
- $Z \approx 1$ dark energy abundance from CMB lensing
- ...

Non-CMB Planck scientific deliverables

- Thousands of galaxy clusters
- Tens of thousands of radio and infrared extra-Galactic sources
- Templates for the diffuse gas in the Galaxy, from 30 to 857 GHz
- ...



Conclusions

- The CMB will be the best signal from the early universe for long
- We have some knowledge of the two point correlation function, but most of the signal is presently unknown
- If detected, the hidden signatures might reveal mysteries for physics, like gravitational waves, or the mechanism of cosmic acceleration
- Planck opening a new era for CMB studies, imaging, all sky polarization, unprecedented foreground control, path finder for even more ambitious mission
- We've no other way to get close to the Big Bang, so let's go for it and see how far we can go
- First go/no go criteria from Planck and other probes in just a few years, possible post-Planck scenarios...



- Polarized foreground too intense, no sufficient cleaning, systematics out of control
- Increase by one digit the cosmological parameters measurement, super-fine knowledge of the Galaxy in the microwave band
- Time scale: few years



Theorist



- Modest controllable foreground emission, systematics under control
- Planck pushes ahead the frontier for particle physics, cosmology and astrophysics
- Cosmological gravity waves discovered from CMB B modes! Expected precision down to one thousandth of the scalar amplitude
- Percent measurement of the dark energy abundance at the onset of acceleration, from CMB lensing
- Time scale: from a few to 30 years



Theorist

Cosmological tensors

Spacetime