



gauge invariance of heat and charge transport coefficients in electronic insulators


Stefano Baroni
Scuola Internazionale Superiore di Studi Avanzati
Trieste — Italy

serious answers to silly questions

- how is it that the heat conductivity is well defined, when the energy current that determines it is not?
- how is it that the electric conductivity of non-ionic fluids vanishes, when the current fluctuations that determine it do not?



conserved extensive quantities (energy, charge, mass, ...) in a macroscopic body flow from high- to low-density regions so as to maximise entropy



the linear-response theory of transport

$$\mathbf{J} = \lambda \nabla \alpha$$

energy transport

$$\alpha = \frac{1}{T}$$

$$J_{\mathcal{E}} = -\kappa \nabla T$$

charge transport

$$\alpha = -\frac{\phi}{T}$$

$$J_Q = \sigma E$$



the linear-response theory of transport

$$\mathbf{J} = \lambda \nabla \alpha$$




Green-Kubo

$$\lambda = \frac{\Omega}{k_B T} \underbrace{\int_0^\infty \langle J(t) J(0) \rangle dt}_{\langle J^2 \rangle \tau}$$




Einstein-Helfand

$$\lambda = \frac{\Omega}{2k_B T} \lim_{t \rightarrow \infty} \frac{1}{t} \text{var} \left[\int_0^t J(t') dt' \right]$$



classical and quantum adiabatic heat transport

$$\mathbf{J}_\varepsilon = \sum_I e_I \mathbf{v}_I + \frac{1}{2} \sum_{I \neq J} (\mathbf{v}_I \cdot \mathbf{F}_{IJ}) (\mathbf{R}_I - \mathbf{R}_J)$$


PRL **104**, 208501 (2010)

PHYSICAL REVIEW LETTERS

week ending
21 MAY 2010

Thermal Conductivity of Periclase (MgO) from First Principles

Stephen Stackhouse*

Department of Geological Sciences, University of Michigan, Ann Arbor, Michigan, 48109-1005, USA

Lars Stixrude[†]

Department of Earth Sciences, University College London, Gower Street, London WC1E 6BT, United Kingdom

Bijaya B. Karki[‡]

*Department of Computer Science, Louisiana State University, Baton Rouge, Louisiana 70803, USA
and Department of Geology and Geophysics, Louisiana State University, Baton Rouge, Louisiana 70803, USA*

sensitive to the form of the potential. The widely used Green-Kubo relation [14] does not serve our purposes, because in first-principles calculations it is impossible to uniquely decompose the total energy into individual contributions from each atom.



how come?



how is it that a formally exact theory of the electronic ground state cannot predict *all* measurable adiabatic properties?

insights from classical mechanics

$$E = \sum_I \epsilon_I(\mathbf{R}, \mathbf{V})$$
$$= \text{cnst}$$

$$\epsilon_I(\mathbf{R}, \mathbf{V}) = \frac{1}{2} M_I V_I^2 + \frac{1}{2} \sum_{J \neq I} v(|\mathbf{R}_I - \mathbf{R}_J|)$$

$$\mathbf{J}_e = \sum_I \epsilon_I \mathbf{V}_I + \frac{1}{2} \sum_{I \neq J} (\mathbf{V}_I \cdot \mathbf{F}_{IJ})(\mathbf{R}_I - \mathbf{R}_J)$$



insights from classical mechanics

$$E = \sum_I \epsilon_I(\mathbf{R}, \mathbf{V})$$

$$\epsilon_I(\mathbf{R}, \mathbf{V}) = \frac{1}{2} M_I V_I^2 + \frac{1}{2} \sum_{J \neq I} v(|\mathbf{R}_I - \mathbf{R}_J|) (1 + \Gamma_{IJ})$$



$$\begin{aligned} \mathbf{J}_e = & \sum_I \epsilon_I \mathbf{V}_I + \frac{1}{2} \sum_{I \neq J} (\mathbf{V}_I \cdot \mathbf{F}_{IJ}) (\mathbf{R}_I - \mathbf{R}_J) \\ & + \frac{1}{2} \sum_{I \neq J} \Gamma_{IJ} [\mathbf{V}_I v(|\mathbf{R}_I - \mathbf{R}_J|) + (\mathbf{V}_I \cdot \mathbf{F}_{IJ}) (\mathbf{R}_I - \mathbf{R}_J)] \end{aligned}$$



insights from classical mechanics

$$\mathbf{J}_e = \sum_I \epsilon_I \mathbf{v}_I + \frac{1}{2} \sum_{I \neq J} (\mathbf{v}_I \cdot \mathbf{F}_{IJ}) (\mathbf{R}_I - \mathbf{R}_J)$$

$$+ \frac{1}{2} \sum_{I \neq J} \Gamma_{IJ} [\mathbf{v}_I v(|\mathbf{R}_I - \mathbf{R}_J|) + (\mathbf{v}_I \cdot \mathbf{F}_{IJ}) (\mathbf{R}_I - \mathbf{R}_J)]$$



insights from classical mechanics

$$\mathbf{J}_e = \sum_I \epsilon_I \mathbf{v}_I + \frac{1}{2} \sum_{I \neq J} (\mathbf{v}_I \cdot \mathbf{F}_{IJ}) (\mathbf{R}_I - \mathbf{R}_J) \\ + \frac{1}{2} \sum_{I \neq J} \Gamma_{IJ} [\mathbf{v}_I v(|\mathbf{R}_I - \mathbf{R}_J|) + (\mathbf{v}_I \cdot \mathbf{F}_{IJ}) (\mathbf{R}_I - \mathbf{R}_J)]$$

$$\dot{\mathbf{P}} = \frac{d}{dt} \frac{1}{4} \sum_{I \neq J} \Gamma_{IJ} v(|\mathbf{R}_I - \mathbf{R}_J|) (\mathbf{R}_I - \mathbf{R}_J)$$



insights from classical mechanics

$$\mathbf{J}' = \mathbf{J} + \dot{\mathbf{P}}$$

$$\kappa \sim \frac{1}{2t} \text{var}[\mathbf{D}(t)] \quad \mathbf{D}(t) = \int_0^t \mathbf{J}(t') dt'$$

$$\mathbf{D}'(t) = \mathbf{D}(t) + \mathbf{P}(t) - \mathbf{P}(0)$$

$$\text{var}[\mathbf{D}'(t)] = \underbrace{\text{var}[\mathbf{D}(t)]}_{\mathcal{O}(t)} + \underbrace{\cancel{\text{var}[\Delta\mathbf{P}(t)]}}_{\mathcal{O}(1)} + \underbrace{\cancel{2\text{cov}[\mathbf{D}(t) \cdot \Delta\mathbf{P}(t)]}}_{\mathcal{O}(t^{\frac{1}{2}})}$$



insights from classical mechanics

$$\mathbf{J}' = \mathbf{J} + \dot{\mathbf{P}}$$


$$\kappa \sim \frac{1}{2t} \text{var} [\mathbf{D}(t)] \quad \mathbf{D}(t) = \int_0^t \mathbf{J}(t') dt'$$

$$\mathbf{D}(t) = \mathbf{D}'(t) + \mathbf{P}(t) - \mathbf{P}(0)$$

$$\kappa' = \kappa$$

$$\text{var} [\mathbf{D}'(t)] = \text{var} [\mathbf{D}(t)] + \text{var} [\mathbf{P}(t)] + 2\text{cov} [\mathbf{D}, \mathbf{P}(t)]$$

gauge invariance of transport coefficients



$$E[\Omega_1 \cup \Omega_2] = E[\Omega_1] + E[\Omega_2]$$

extensivity

$$\mathcal{E}[\Omega] = \int_{\Omega} e(\mathbf{r}) d\mathbf{r}$$

$$\mathbf{J}(t) = \frac{1}{\Omega} \int \mathbf{j}(\mathbf{r}, t) d\mathbf{r}$$

thermodynamic invariance

$$\mathcal{E}'[\Omega] = \mathcal{E}[\Omega] + \mathcal{O}[\partial\Omega]$$

$$\mathbf{P}(t) = \frac{1}{\Omega} \int \mathbf{p}(\mathbf{r}, t) d\mathbf{r}$$

gauge invariance

$$e'(\mathbf{r}) = e(\mathbf{r}) - \nabla \cdot \mathbf{p}(\mathbf{r})$$

$$\mathbf{j}'(\mathbf{r}, t) = \mathbf{j}(\mathbf{r}, t) + \dot{\mathbf{p}}(\mathbf{r}, t)$$

$$\mathbf{J}'(t) = \mathbf{J}(t) + \dot{\mathbf{P}}(t)$$

conservation

$$\dot{e}(\mathbf{r}, t) = -\nabla \cdot \mathbf{j}(\mathbf{r}, t)$$

gauge invariance of transport coefficients

$$E[\Omega_1 \cup \Omega_2] = E[\Omega_1] + E[\Omega_2]$$

any two conserved densities that differ by the divergence of a (bounded) vector field are physically equivalent

the corresponding conserved fluxes differ by a total time derivative, and the transport coefficients coincide

thermodynamic invariance

$$\mathcal{E}'[\Omega] = \mathcal{E}[\Omega] + \mathcal{O}[\partial\Omega]$$

$$P(t) = \frac{1}{\Omega} \int p(\mathbf{r}, t) d\mathbf{r}$$

gauge invariance

nature
physics

ARTICLES

PUBLISHED ONLINE: 19 OCTOBER 2015 | DOI: 10.1038/NPHYS3509

Microscopic theory and quantum simulation of atomic heat transport

Aris Marcolongo¹, Paolo Umari² and Stefano Baroni^{1*}

$$J'(t) = J(t) + \dot{P}(t)$$

conservation



gauge invariance of heat transport

$$\mathbf{J}_\varepsilon = \sum_I e_I \mathbf{v}_I + \frac{1}{2} \sum_{I \neq J} (\mathbf{v}_I \cdot \mathbf{F}_{IJ}) (\mathbf{R}_I - \mathbf{R}_J)$$

PRL **104**, 208501 (2010)

PHYSICAL REVIEW LETTERS

week ending
21 MAY 2010

Thermal Conductivity of Periclase (MgO) from First Principles

Stephen Stackhouse*

Department of Geological Sciences, University of Michigan, Ann Arbor, Michigan, 48109-1005, USA

Lars Stixrude†

Department of Earth Sciences, University College London, Gower Street, London WC1E 6BT, United Kingdom

Bijaya B. Karki‡

*Department of Computer Science, Louisiana State University, Baton Rouge, Louisiana 70803, USA
and Department of Geology and Geophysics, Louisiana State University, Baton Rouge, Louisiana 70803, USA*





sensitive to the form of the potential. The widely used Green-Kubo relation [14] does not serve our purposes, because in first-principles calculations it is impossible to uniquely decompose the total energy into individual contributions from each atom.

solution:

choose *any* local representation of the energy that integrates to the correct value and whose correlations decay at large distance — the conductivity computed from the resulting current will be *independent* of the chosen representation.



thermal conductivity of liquid water from DFT




breaking news: $\kappa_{SCAN} \approx 0.8 \left[\frac{W}{mK} \right]$

ionic transport





+
+
+





-
-
-

$$\mathbf{J} = \sigma \mathbf{E}$$

$$\mathbf{J} = \sum_i q_i \mathbf{v}_i$$

$$\sigma = \frac{\Omega}{3k_B T} \langle |\mathbf{J}|^2 \rangle \times \tau_J$$







$$\mathbf{J} = \sigma \mathbf{E}$$

$$\mathbf{J} = ???$$

$$\sigma = \frac{\Omega}{3k_B T} \langle |\mathbf{J}|^2 \rangle \times \tau_J$$






$$\mathbf{J} = \sigma \mathbf{E}$$

$$\begin{aligned} \mathbf{J} &= \frac{1}{\Omega} \dot{\boldsymbol{\mu}} \\ &= \frac{1}{\Omega} \sum_i \mathbf{z}_i^* \cdot \mathbf{v}_i \end{aligned}$$

$z_{i\alpha\beta}^* = \frac{\partial \mu_\alpha}{\partial u_{i\beta}}$

$$\sigma = \frac{\Omega}{3k_B T} \langle |\mathbf{J}|^2 \rangle \times \tau_J$$

the conundrum



pure, undissociated
 H_2O

$$\mathbf{J} = \frac{1}{\Omega} \sum_i \mathbf{z}_i^* \cdot \mathbf{v}_i$$

$$\neq 0$$

$$\sigma = \frac{\Omega}{3k_B T} \langle |\mathbf{J}|^2 \rangle \times \tau_J$$

$$= 0$$

???

the conundrum

PRL 107, 185901 (2011)

PHYSICAL REVIEW LETTERS

week ending
28 OCTOBER 2011

Dynamical Screening and Ionic Conductivity in Water from *Ab Initio* Simulations

Martin French,¹ Sebastien Hamel,² and Ronald Redmer¹

“Interestingly, the use of predefined constant charges can yield the same conductivity as is found with the fully time-dependent charge tensors, but only if they have values of $Z_H=1$ and $Z_O=-2$.”


atomic
“oxidation states”





how come?

gauge invariance of charge transport





$$\sigma \propto \lim_{t \rightarrow \infty} \frac{1}{2t} \text{var} [\mu_{AB}(t)]$$

$$\mu_{AB}(t) = \int_0^t J(t') dt'$$



$$\text{var} [\mu_{AB}] = \underbrace{\text{var} [\mu_{AA'}]}_{\mathcal{O}(t)} + \underbrace{\text{var} [\mu_{A'B}]}_{\mathcal{O}(1)} + 2 \underbrace{\text{cov} [\mu_{AA'} \cdot \mu_{A'B}]}_{\mathcal{O}(t^{\frac{1}{2}})}$$

$$\sigma \propto \lim_{t \rightarrow \infty} \frac{1}{2t} \langle |\mu_{AA'}|^2 \rangle$$



topological invariants



topological invariants





topological invariants






$$Q(AA') = Q(AA') = Q[n_1 = 1, n_2 = 1]$$

a numerical experiment on molten KCl




a topologically non-trivial minimum-energy path
connecting two identical configurations of a ionic fluid

a numerical experiment on molten KCl




effective charge





topological charge

$$q_x = -0.000(6); \quad q_y = 0.000(2); \quad q_z = 1.00(18)$$

a numerical experiment on molten KCl



$$\begin{aligned} Q_z[\text{Cl}] &= -1 & Q_y[\text{Cl}] &= -1 \\ Q_z[\text{K}] &= 1 & Q_z[\text{K}] &= 0 \end{aligned}$$




the charges transported by K and Cl
around z cancel exactly

atomic oxidation states

$$Q_\alpha[\mathcal{C}] = \frac{1}{\ell} \mu_\alpha[\mathcal{C}]$$

$$Q_\alpha[\mathcal{C}_1 \circ \mathcal{C}_2] = Q_\alpha[\mathcal{C}_1] + Q_\alpha[\mathcal{C}_2]$$

- All loops can be shrunk to a point without closing the gap (*strong adiabaticity*);
- Any two like atoms can be swapped without closing the gap



$$q_{i\alpha\beta} = q_{S(i)} \delta_{\alpha\beta}$$

atomic oxidation state

currents from atomic oxidation numbers

$$J_{\alpha} = \sum_{i\beta} Z_{i\alpha\beta}^* V_{i\beta} \quad (2)$$

$$J'_{\alpha} = \sum_i q_{S(i)} V_{i\alpha} \quad (9)$$


nature
physics

ARTICLES

<https://doi.org/10.1038/s41567-019-0562-0>

Topological quantization and gauge invariance of charge transport in liquid insulators







Federico Grasselli¹ and Stefano Baroni^{1,2*}




$$\left\langle \left| \int_0^t \mathbf{J}(t') dt' \right|^2 \right\rangle$$



non-stoichiometric melts






breach of strong adiabaticity




$$\mu \neq \mu^*$$

$$\mu \neq 0$$

weakly adiabatic transport



weakly adiabatic transport



$$\Delta\mu = e \int_0^t \mathbf{J}(t') dt'$$

$$J_\alpha(t) = \sum_{i\beta} Z_{i\alpha\beta}^*(t) v_{i\beta}(t)$$

$$J_\alpha(t) = \sum_i q_{S(i)} v_{i\alpha}(t) - 2v_\alpha^{lp}(t)$$

cross term


PHYSICAL REVIEW X

Oxidation States, Thouless' Pumps, and Nontrivial Ionic Transport in Nonstoichiometric Electrolytes

Paolo Pegolo, Federico Grasselli, and Stefano Baroni
Phys. Rev. X **10**, 041031 – Published 12 November 2020




strongly adiabatic transport



$$\begin{aligned} \mu &= \mu^* \\ \mu &= 0 \end{aligned}$$



weakly adiabatic transport



$$\begin{aligned} \mu &\neq \mu^* \\ \mu &\neq 0 \end{aligned}$$



conclusions

- conserved currents are intrinsically ill-defined at the atomic scale;
- conservation and extensiveness make transport coefficients independent of the specific microscopic representation of the conserved densities and currents;
- this *gauge invariance* of transport coefficients makes it possible to compute thermal transport coefficients from DFT using equilibrium AIMD and the Green-Kubo formalism;
- in the (quasi-) harmonic approximation, the theory smoothly interpolates between the BTE in crystals and the Allen-Feldman model of heat conduction in glasses;
- topological quantisation of charge transport allows one to give a rigorous definition of the atomic oxidation states;
- gauge invariance and topological quantisation of charge transport make the electric conductivity of stoichiometry ionic conductors depend on the formal oxidation numbers of the ionic species, via the Green-Kubo formula;
- in non-stoichiometric ionic conductors, the breach of the topological conditions that allow a univocal definition of the atomic oxidation numbers determine a regime where charge can flow without any associated mass drift.



supported by:



<http://www.quantum-espresso.org>



<http://foundation.quantum-espresso.org>



<http://www.max-centre.eu>





thanks to:



Aris Marcolongo



Loris Ercole



Riccardo Bertossa



Leyla Isaeva



Giuseppe Barbalinardo



Davide Donadio



Federico Grasselli



Paolo Pegolo



Davide Tisi

nature physics

ARTICLES

PUBLISHED ONLINE: 19 OCTOBER 2015 | DOI: 10.1038/NPHYS3509

Microscopic theory and quantum simulation of atomic heat transport

Aris Marcolongo¹, Paolo Umari² and Stefano Baroni^{1*}


nature physics

ARTICLES

<https://doi.org/10.1038/s41567-019-0562-0>

Topological quantization and gauge invariance of charge transport in liquid insulators

Federico Grasselli¹ and Stefano Baroni^{1,2*}



ARTICLE

<https://doi.org/10.1038/s41467-019-11572-4>

OPEN

Modeling heat transport in crystals and glasses from a unified lattice-dynamical approach

Leyla Isaeva¹, Giuseppe Barbalinardo², Davide Donadio² & Stefano Baroni^{1,3}

PHYSICAL REVIEW X

Oxidation States, Thouless' Pumps, and Nontrivial Ionic Transport in Nonstoichiometric Electrolytes

Paolo Pegolo, Federico Grasselli, and Stefano Baroni
Phys. Rev. X **10**, 041031 – Published 12 November 2020