

The exact theory of phase separation and interfaces in near-critical planar systems has been introduced and developed in the papers [1-7]. The derivation in three dimensions has been obtained in [8].

[1] DELFINO G, VITI J (2012). Phase separation and interface structure in two dimensions from field theory. JOURNAL OF STATISTICAL MECHANICS: THEORY AND EXPERIMENT, P10009, doi: 10.1088/1742-5468/2012/10/P10009

[2] Gesualdo Delfino, Alessio Squarcini,
Interfaces and wetting transition on the half plane. Exact results from field theory,
J. Stat. Mech. 2013 (2013) P05010,
DOI: 10.1088/1742-5468/2013/05/P05010

[3] Gesualdo Delfino and Alessio Squarcini,
Exact theory of intermediate phases in two dimensions,
Annals of Physics 342 (2014) 171,
DOI: 10.1016/j.aop.2013.12.010

[4] Gesualdo Delfino and Alessio Squarcini,
Phase separation in a wedge. Exact results,
Phys. Rev. Lett. 113 (2014) 066101,
DOI: <http://dx.doi.org/10.1103/PhysRevLett.113.066101>

[5] Gesualdo Delfino and Alessio Squarcini,
Multiple phases and vicious walkers in a wedge,
Nucl. Phys. B 901 (2015) 430-443
doi: 10.1016/j.nuclphysb.2015.10.019

[6] Gesualdo Delfino,
Interface localization near criticality,
JHEP 05 (2016) 032
doi: 10.1007/JHEP05(2016)032

[7] Gesualdo Delfino and Alessio Squarcini,
Long range correlations generated by phase separation. Exact results from field theory,
JHEP 11 (2016) 119
doi: 10.1007/JHEP11(2016)119

[8] Gesualdo Delfino, Walter Selke and Alessio Squarcini,
Particles, string and interface in the three-dimensional Ising model,
Nucl. Phys. B 958 (2020) 115139
doi: 10.1016/j.nuclphysb.2020.115139